

Demanding diagnostics – important diagnosis

EDUCATIONAL CASE REPORT

ØIVIND EKEBERG

Øivind Ekeberg (born 1945) Specialist in psychiatry and supervisor in psychodynamic psychotherapy. Senior Consultant in psychiatry, Department of Acute Medicine, Oslo University Hospital, Ullevål. Adjunct Professor in medical behavioural science, Institute of Basal Medical Sciences, University of Oslo. He works as both clinician and researcher, particularly with suicidal behaviour, psychotraumatology and mental reactions in connection with somatic disease.

The author has completed the ICMJE form and declares no conflicts of interest.

Email: oivind.ekeberg@ous-hf.no
Oslo University Hospital, Ullevål

There is little probability of a healthy patient who has not previously had psychiatric problems becoming manic in her late 50s without there being an organic reason for it.

The somatic diagnostics may be demanding. It is necessary to examine possible neurological, metabolic, medicinal and intoxicant-related causes. The patient had early symptoms such as impaired strength and unsteady gait which pointed in a neurological direction. Many conditions do not show up on a CT or, less commonly, on an MRI either. A lumbar puncture should be performed early.

Antibodies develop slowly in borreliosis, and serological tests are frequently negative, particularly in the early stages of the disease (1). The incubation period may be long, so that in many cases the patient is not aware of exposure. The diagnostic challenges can be illustrated by the fact that the disease is also called «the new great imitator». Neuroborreliosis has become the most frequently occurring neuroinfectious condition in Europe where the source of infection is a vector (2). The prevalence is increasing, probably partly due to vegetation changes and a milder climate.

The most common psychiatric symptoms are tiredness and depression, but panic attacks, schizophrenia-like symptoms and hallucinations may also occur [\(2\)](#).

The patient was hospitalised two months later on suspicion of depression, but the diagnosis was not confirmed. However, she was suffering from tiredness, a common sequela that can last a long time. After completion of treatment, regional hypoactivity with respect to blood supply and metabolism was also found, particularly in the temporal and parietal lobes and in the limbic system [\(3\)](#).

In addition to neurological and psychiatric symptoms, skin manifestations in particular are common, but the heart and limbs may also be affected.

Although the prognosis is normally good, a sneaking development may mean that more areas of the brain may be affected before diagnosis and treatment, resulting in a protracted course for the disease.

The treatment may be difficult because the bacterium has developed many defence mechanisms. The treatment is often successful if it is applied early, but a chronic infection may be difficult to treat [\(1\)](#).

Early diagnosis and treatment is important with a view to shortening both the course of the active disease and the restitution period. Since the symptoms presented can be so variable, it is useful for rare cases, such as the present one, to be published.

LITERATURE

1. Brorson Ø. *Borrelia burgdorferi – en unik bakterie*. Tidsskr Nor Legeforen 2009; 129: 2114–7. [PubMed]
2. Bär KJ, Jochum T, Häger F et al. Painful hallucinations and somatic delusions in a patient with the possible diagnosis of neuroborreliosis. *Clin J Pain* 2005; 21: 362 – 3. [PubMed] [CrossRef]
3. Fallon BA, Lipkin RB, Corbera KM et al. Regional cerebral blood flow and metabolic rate in persistent Lyme encephalopathy. *Arch Gen Psychiatry* 2009; 66: 554 – 63. [PubMed] [CrossRef]

Publisert: 6 March 2012. Tidsskr Nor Legeforen. DOI: 10.4045/tidsskr.11.1288
Submitted 30 October 2011 and approved 3 November 2011. Medical editor: Erlend Hem.
© Tidsskrift for Den norske legeforening 2026. Downloaded from tidsskriftet.no 10 February 2026.