
12 years of «Case reports» in the Journal of the Norwegian Medical Association

ORIGINAL ARTICLE

ERLEND T. AASHEIM

Erlend T. Aasheim (born 1974) MD PhD, is medical editor of the Journal of the Norwegian Medical Association and Academic Clinical Fellow at the Department of Public Health and Primary Care, University of Cambridge, England.

Conflicts of interest: None declared.

Email: erlend.aasheim@legeforeningen.no

Journal of the Norwegian Medical Association
and

Department of Public Health and Primary Care
University of Cambridge
England

Background.

«Case reports» is the title of a column in the Journal of the Norwegian Medical Association devoted to educational patient stories accompanied by an expert commentary. The aim of this study is to describe the patients and authors in this column since its introduction in 1999.

Material and methods.

The study is based on electronic literature searches in the Journal's internet archive (2000–2011) supplemented by manual searches in the printed edition (1999). The medical specialties of the authors were defined on the basis of their workplace affiliations.

Results.

Literature searches identified 157 case reports with accompanying expert commentaries. Among the 157 patients included there were 84 (54%) women. The most frequent age category was 51–60 years with an age range from 0–90 years. First authors were most often affiliated to departments of internal

medicine with accompanying subspecialties (34%); neurology (13%); general surgery with subspecialties (11%); and paediatric departments (10%). Para-clinical specialties were mainly represented by co-authorships; most frequent were histopathology and radiology. Of 676 authorships, only 7 (1.0%) were affiliated to psychiatry and only 5 (0.7%) to general practice.

Interpretation.

The column «Case reports» in the Journal of the Norwegian Medical Association has in the past 12 years illustrated a wide range of patients, but the medical specialties are unevenly represented. It might be desirable to have more articles originating from psychiatry and general practice.

Anecdotal patient descriptions have been of great importance for the development of clinical medicine and previously had a central place in medical publishing. This tradition is also true for Norway [\(1, 2\)](#). But does the anecdotal patient history still have a place in today's evidence-based medicine?

Randomised studies and meta-analyses provide the best basis for clinical treatment guidelines. This means that case reports now play a smaller role in medical publishing. But there are several good reasons why case reports should continue to be published. The thalidomide story showed how individual observations could change clinical practice [\(3\)](#). Among the drugs withdrawn from the market in Spain from 1990 to 1999 in the interests of patient safety, case reports were the most important information source in 18 of 22 instances [\(4\)](#).

Case reports have an intuitive appeal and capture the reader's interest in a different way than a meta-analysis. Case reports also have an important function for doctors in training [\(5\)](#). For these reasons, since 1999 the Journal has published case reports as educational patient stories, accompanied by expert commentaries, in a dedicated column [\(6\)](#). The main point of these case reports is not to present rare conditions, but to promote excellence in clinical medical thinking [\(2\)](#). In a reader survey for the Journal in 2007, many readers stated that they would like to see more case reports [\(7\)](#).

For a general medical journal such as the Journal, case reports should ideally represent a diversity of patients and subjects. Does this happen? Which patients are described, and to what extent are the various medical specialties represented? When the column was introduced in 1999, the vision was that it would represent a relay race from hospital to hospital across the country, also outside the university hospitals [\(6\)](#). To what extent has this goal been achieved? This study has surveyed patients and authors for the 157 case reports and expert commentaries published in the period 1999–2011.

Materials and methods


Relevant articles were identified on *www.tidsskriftet.no* 31.12. 2011 with the search term «Noe å lære av» (the Norwegian name of the case report column) and the advanced search feature «With the exact phrase». The search identified 339 articles published from 10 June 2000 to 13 December 2011. Articles were excluded if they were published in other columns (n = 42), if they were English translations (n = 6) or referred to a larger group of patients (n = 1). The remaining articles were Norwegian-language case reports (n = 145) with accompanying commentaries (n = 145). Manual searches in the journal's print edition for 1999 identified an additional 12 relevant case reports with accompanying commentaries.

For the patients described in the articles, data were recorded on primary diagnosis, gender, pregnancy and age category (0–1 years, 2–10 years, 11–20 years, 21–30 years etc.). For articles that described several patients with the same primary diagnosis (n = 2), only data on the first patient were used.

For the authors, all workplace related affiliations were recorded. Each author was then classified by geographical location (1 of 19 counties) and medical specialty (1 of 44 specialties). Classification of specialty was made based on the author's primary place of work. When workplace indicated a sub-specialty (e.g., «Cardiac Department»), the sub-specialty was recorded and not the main specialty. If in doubt when assigning specialty, information about secondary work place was used. If still in doubt, a preliminary search on the author's name in Google was done, followed by targeted searches among accredited specialists on the Norwegian Medical Association's website *www.legeforeningen.no*. If still in doubt, the author was classified under «unknown specialty» (n = 4). For workplaces with the words «emergency medicine» in the name, searches were made on the author's name among accredited specialists in anaesthesiology, cardiology, internal medicine and psychiatry; writers who were not listed in any of these were classified under anaesthetics. Two authors were classified as psychologists, two as clinical scientists, one as a dentist and one as a pharmacist.

Results

Literature searches identified a total of 157 case reports with 157 accompanying expert commentaries published in the study period. After the introduction of the column in April 1999, 12 case reports were published in the same year. The number then fell to 0 case reports in 2003, followed by an increase to a peak of 23 case reports in the years 2007 and 2008 (e-fig. 1).



e-figure 1 Case reports and commentaries in the Journal of the Norwegian Medical Association from 20 April 1999 to 31 December 2011

Of the 157 patients, 54% were women. The most common age category was 51–60 years. Among the women, 13 pregnancies were noted. Most of the main diagnoses recorded were subject to only one case report, with the exception of three articles about endocarditis and two papers each for amyloidosis, dermatomyositis, Kawasaki disease, leishmaniasis, epidemic nephropathy, and tuberculosis.

The articles were written by a total of 676 authors, representing 157 first authors and 359 co-authors of case reports, and 160 authors of the accompanying commentaries. Case reports had a median of 3 authors (range 1–7). All commentaries, except for three with two authors, had a single author.

Medical specialties

The 157 first authors most often worked at departments of internal medicine and its sub-specialties ($n = 54$). This was followed by neurology ($n = 21$), general surgery with sub-specialties ($n = 18$), paediatrics ($n = 15$) and anaesthetics ($n = 10$). The non-clinical specialties were less often represented, and mainly by co-authorships. Most frequently represented were pathology ($n = 27$) and radiology ($n = 22$).

When related to the number of accredited specialists (8), neurology had most authorships (Fig. 2). The large specialties general medicine and psychiatry were poorly represented and contributed 5 and 7 of 676 authorships, respectively.




Figure 2 Medical specialty of 676 authors in the column «Case reports» with accompanying commentaries shown with the total number of approved specialists in Norway (8)

Counties

Most first authorships originated from Oslo (n = 47), Nordland (n = 17) and Hordaland (n = 15). When related to the number of physicians in each county (9), Nordland, Sogn og Fjordane and Troms had most first authorships (Fig. 3). Accompanying commentaries (invited by the editors) most often originated from Bergen, Oslo and Sør-Trøndelag, both with and without adjustment for the number of doctors in each county.




Figure 3 County affiliation of 676 authors of case reports and commentaries, shown with the total number of doctors in each county (9)

Discussion

This study shows that case reports and accompanying expert commentaries during a period of 12 years have presented a broad range of patients and diagnoses, whilst the medical specialties have been unevenly represented. Domination of neurological cases is consistent with the pattern for case reports published in *The Lancet* (10). A majority of first authors from clinical specialties is not unexpected for case reports, but remarkably few case reports originated from psychiatry and general practice. Primary care has previously been encouraged to contribute to this column (2), but the presented findings provide a rationale for also extending this invitation to other specialties.

Author lists that include five or six different specialties (11,12) suggest impressive multi-disciplinary collaboration. Probably all clinicians who have been actively involved in the diagnostic work-up and treatment of a patient should be informed before a case report is submitted for publication. It has been pointed out that radiologists and other specialties are not always represented in the author list when a case report is published, even if they have contributed substantially in the diagnostic process and treatment of the patient (13).

The number of authorships in the counties varied, but all counties were represented. This is particularly interesting, considering that the original scheme of inviting contributions to the column (6) rapidly collapsed (2). It is worth noting that individual efforts impacted on the specialty and county level: Erik Waage Nielsen (anaesthetist in Bodø) and Odd Kildahl Andersen (internal medicine specialist in Harstad) have authored 11 and seven case reports in the column, respectively. These efforts are impressive at a time when case reports may yield less academic merit than original articles.

The study has several limitations. Authors with affiliation to a given specialty, e.g. general practice, may have written a case report while working in another specialty, e.g. internal medicine, and if so, this would not have been identified. The great majority of manuscripts were submitted spontaneously throughout most of the study period, while expert commentaries were invited by the editors. Information about rejected manuscripts was not available. It is therefore not known whether the published case reports are representative of all case report manuscripts submitted for consideration for publication.

Tabell

Main points
<ul style="list-style-type: none">• The «Case reports» column contains educational patient stories accompanied by expert commentaries• A review of the column shows that a wide range of patients are described• More contributions from general medicine and psychiatry could add valuable perspectives

LITERATURE

1. Hem E. Enkeltpasienter som læremestre. Tidsskr Nor Lægeforen 2007; 127: 561. [PubMed]
2. Wyller TB. Kasuistikker for det 21. århundre. Tidsskr Nor Lægeforen 2004; 124: 1913. [PubMed]
3. Mcbride WG. Thalidomide and congenital abnormalities. Lancet 1961; 278: 1358. [CrossRef]
4. Arnaiz JA, Carné X, Riba N et al. The use of evidence in pharmacovigilance. Case reports as the reference source for drug withdrawals. Eur J Clin Pharmacol 2001; 57: 89 – 91. [PubMed] [CrossRef]
5. Vandenbroucke JP. In defense of case reports and case series. Ann Intern Med 2001; 134: 330 – 4. [PubMed]
6. Gulbrandsen P. Noe vi lærte av er noe å skrive om. Tidsskr Nor Lægeforen 1999; 119: 1265 – 6. [PubMed]

7. Aasland OG. Tidsskriftets leserundersøkelse 2007. Tidsskr Nor Legeforen 2008; 128: 1512-4. [PubMed]
 8. Den norske legeforening. Godkjente spesialister per 13.4.2011. www.legeforeningen.no/id/171362 (6.1.2012).
 9. Den norske legeforening. Alle legemedlemmer fordelt på fylkesavdeling og yrkesforening per 3.9.2007. www.legeforeningen.no/id/99171 (6.1.2012).
 10. Kang S. Anecdotes in medicine – 15 years of Lancet case reports. Lancet 2010; 376: 1448 – 9. [PubMed] [CrossRef]
 11. Zätterstrøm UK, Hokland BM, Walberg M et al. En halsinfeksjon med rask forverring. Tidsskr Nor Lægeforen 2002; 122: 1896 – 9. [PubMed]
 12. Moser KH, Rødevand E, Hammerstrøm J et al. En tidligere frisk mann med feberepisoder med kirkespirforløp. Tidsskr Nor Legeforen 2008; 128: 1845-7. [PubMed]
 13. Stiris MG. Fortsatt noe å lære av? Tidsskr Nor Lægeforen 2007; 127: 1234. [PubMed]
-

Publisert: 27 March 2012. Tidsskr Nor Legeforen. DOI: 10.4045/tidsskr.11.1264

Received 24 October 2011, first revision submitted 13 November 2011, approved 1 December 2011. Medical editor Petter Gjersvik.

© Tidsskrift for Den norske legeforening 2026. Downloaded from tidsskriftet.no 28 March 2026.