

---

# Artery of Percheron occlusion

---

## IMAGES IN MEDICINE

ARNHILD ØSTERÅS BØGSETH

Section of Medicine

Lovisenberg Diaconal Hospital

Arnhild Østerås Bøgseth, specialty registrar in internal medicine.

The author has completed the ICMJE form and declares no conflicts of interest.

JELENA ZUGIC SOARES

E-mail: [jelenazugic.soares@lds.no](mailto:jelenazugic.soares@lds.no)

Unger-Vetlesen Institute

Lovisenberg Diaconal Hospital

Jelena Zugic Soares, specialist in internal medicine and PhD candidate.

The author has completed the ICMJE form and declares no conflicts of interest.

RAGNHILD UNDSETH

Section of Surgery

Lovisenberg Diaconal Hospital

Ragnhild Undseth, PhD, senior consultant and specialist in radiology.

The author has completed the ICMJE form and declares no conflicts of interest.

JØRGEN VALEUR


Unger-Vetlesen Institute

Lovisenberg Diaconal Hospital

Jørgen Valeur, PhD, specialty registrar in internal medicine and gastroenterology, and head of department.

The author has completed the ICMJE form and declares no conflicts of interest.

Arnhild Østerås Bøgseth and Jelena Zugic Soares have made an equal contribution to this article.



The thalamus is normally supplied with blood on both sides by small branches leading from the ipsilateral posterior communicating artery, as well as P1 and P2 segments of the posterior cerebral artery. The most frequent cause of injury in the thalamus is thrombosis in parts of the posterior cerebral artery. In 1973, the French neurologist Gérard Percheron (1930–2011) described a rare anatomical variant in which a single trunk from the posterior cerebral artery supplies medial parts of the thalamus and ventral parts of the mesencephalon bilaterally [\(1\)](#). This anatomical variant is called the artery of Percheron (illustration on left).

Percheron artery occlusion results in bilateral thalamic infarcts, a rare condition. It is rarely possible to detect arterial occlusion with diagnostic imaging, and diagnosis is made indirectly through findings of bilateral infarcts in the medial thalamus, with or without affection of the mesencephalon [\(2\)](#).

The symptoms can be dramatic, with acute onset of coma in addition to focal neurological, neuropsychological and ophthalmological signs (abnormal pupillary light reflexes, reduced ocular motility, or ptosis). Early diagnosis can be challenging, as the clinical disease picture is unusual and radiological examinations in the initial phase may not yield definitive pathological findings [\(3\)](#).

Our patient was a man in his late sixties who abruptly lost consciousness while on holiday abroad. He was quickly collected by ambulance and driven to the nearest hospital. Thrombolytic therapy was not initiated because the cause of his loss of consciousness was unclear. He was intubated on arrival and the left pupil was observed to be more dilated than the right one. CT head and CT angiography performed on the day of admission yielded no relevant pathological findings. Head MRI on day 4 of the hospitalisation showed infarcts bilaterally in the medial thalamus (image on right).

The patient was extubated on day 5 and transferred to a local hospital in Norway 14 days later. He had partial tetraplegia with reduced function in the left-sided extremities and paralysis in the right-sided extremities. He was unable to open his eyes, but his vision was intact if the eyelids were lifted for him. His level of consciousness fluctuated, but gradually improved. His swallowing function gradually returned but did not normalise completely. A PEG tube was therefore inserted. With physiotherapy he eventually managed to

walk 50 m using a walking frame. The patient's functional level was too low to benefit from the rehabilitation department, and he was placed permanently in a nursing home. The disease course was complicated with recurrent aspiration pneumonia, and he died approximately one year later.

---

*The patient's next of kin have consented to the publication of this article.*

---

## LITERATURE

1. Percheron G. The anatomy of the arterial supply of the human thalamus and its use for the interpretation of the thalamic vascular pathology. *Z Neurol* 1973; 205: 1 - 13. [PubMed]
  2. Arauz A, Patiño-Rodríguez HM, Vargas-González JC et al. Clinical spectrum of artery of Percheron infarct: clinical-radiological correlations. *J Stroke Cerebrovasc Dis* 2014; 23: 1083 - 8. [PubMed][CrossRef]
  3. Stamm BJ, Lineback CM, Skolarus LE et al. Artery of Percheron infarct: 12 cases and their complex clinical courses. *Neurohospitalist* 2018; 8: 141 - 5. [PubMed][CrossRef]
- 

Publisert: 15 November 2018. *Tidsskr Nor Legeforen*. DOI: 10.4045/tidsskr.18.0289  
Received 5.4.2018, first revision submitted 14.8.2018, accepted 17.9.2018.  
Copyright: © Tidsskriftet 2026 Downloaded from tidsskriftet.no 29 March 2026.