

---

# Saving lives together

---

## PERSPECTIVES

CONRAD ARNFINN BJØRSHOL

E-mail: [conrad.bjorshol@sus.no](mailto:conrad.bjorshol@sus.no)

Conrad Arnfinn Bjørshol, chief medical advisor for the *Sammen redder vi liv* campaign, senior researcher at the Regional Centre for Emergency Medical Research and Development (RAKOS), Western Norway Health Authority, and senior consultant at the Department of Anaesthesiology and Intensive Care, Stavanger University Hospital.

The author has completed the ICMJE form and declares the following conflict of interest: He has participated in meetings of the Global Resuscitation Alliance with financial support from the Laerdal Foundation for Acute Medicine, TrygFonden and EMS 2018 (European Emergency Medical Services Congress).

BJØRN JAMTLI

Bjørn Jamtli, senior advisor in the Department of Emergency Medicine and Preparedness, Norwegian Directorate of Health.

The author has completed the ICMJE form and declares no conflicts of interest.

JO KRAMER-JOHANSEN

Jo Kramer-Johansen, PhD, anaesthesiologist at the Air Ambulance Department, Oslo University Hospital, and professor II at the Norwegian National Advisory Unit on Prehospital Emergency Medicine (NAKOS) and Institute of Clinical Medicine, University of Oslo. He is clinical director of the Norwegian Cardiac Arrest Registry and the Norwegian AED Registry.

The author has completed the ICMJE form and declares no conflicts of interest.

HELGE MYKLEBUST

Helge Myklebust, research director of Laerdal Medical, has assisted the Directorate of Health in the *Sammen redder vi liv* campaign.

The author has completed the ICMJE form and declares no conflicts of interest.

#### STEINAR OLSEN

Steinar Olsen, head of the Department of Emergency Medicine and Preparedness, Norwegian Directorate of Health.

The author has completed the ICMJE form and declares no conflicts of interest

#### TROND NORDSETH

Trond Nordseth, PhD, specialist in anaesthesiology and senior consultant at Molde Hospital and St. Olavs Hospital. He is head of the Norwegian Resuscitation Council (NRR).

The author has completed the ICMJE form and declares no conflicts of interest.

#### SVEN MOLLEKLEIV

Sven Mollekleiv, senior vice president of DNV GL and honorary president of the Norwegian Red Cross.

The author has completed the ICMJE form and declares no conflicts of interest.

#### SIW LILLY OSMUNDSEN

Siw Lilly Osmundsen, senior advisor and head of the Norwegian AED Registry, Norwegian National Advisory Unit on Prehospital Emergency Medicine (NAKOS). She is a board member of the Norwegian First Aid Council.

The author has completed the ICMJE form and declares no conflicts of interest.

#### JAN ERIK NILSEN

Jan Erik Nilsen, senior consultant at the Pre-Hospital Clinic, Oslo University Hospital, and director of the Norwegian National Advisory Unit on Prehospital Emergency Medicine (NAKOS).

The author has completed the ICMJE form and declares no conflicts of interest.

#### HANS MORTEN LOSSIUS

Hans Morten Lossius, PhD, specialist in anaesthesiology and professor of advanced pre-hospital emergency medicine at the University of Stavanger and Aalborg University. He is general secretary of the Norwegian Air Ambulance Foundation.

The author has completed the ICMJE form and declares no conflicts of interest.

---

**Training in first aid has a long tradition in Norway. We believe that survival after time-critical events outside hospital can be further improved through systematic training.**



Illustration: Helene Brox

The Directorate of Health seeks to engage all ready and willing forces in a collaboration aiming to increase the survival rate after cardiac arrest and other time-critical incidents outside hospitals. The reason is that the population is an especially crucial actor in the medical emergency chain [\(1\)](#). Hence, good, life-long learning is called for. The voluntary organisations should continue to play an important role. Programmes that coordinate efforts across multiple links in the medical emergency chain have been shown to improve survival rates, for example in Denmark [\(2\)](#).

---

## Time-critical conditions

Out-of-hospital cardiac arrest (OHCA) is the most time-critical emergency condition of all. Each year, laypeople or ambulance personnel attempt to resuscitate approximately 3 000 persons who have suffered an out-of-hospital cardiac arrest [\(3\)](#). The proportion of patients with cardiac arrest who receive cardiopulmonary resuscitation (CPR) before the ambulance arrives is high in Norway (over 80 %) [\(3\)](#). In 2017, altogether 368 patients survived for at least 30 days after suffering out-of-hospital cardiac arrest in Norway (nearly 14 % of

those who were treated by paramedics). This is equivalent to 6.8 per 100 000 inhabitants, but the figures vary from one health trust to another [\(3\)](#). In Denmark, the average 30-day survival rate amounted to 7.7 per 100 000 in the years 2012–16 [\(4\)](#). If Norway could achieve the same survival rate as Denmark, 47 more people would survive each year.

Injuries and poisonings cause approximately 2 500 deaths each year in Norway, of which approximately 600 are suicides. With an annual incidence of 40 per 100 000 inhabitants, serious injuries account for approximately 6 % of all deaths in Norway. When it comes to serious injuries, it is estimated that 1.8–4.5 % of those who die may be saved if bystanders help keep the airways open and stop major haemorrhages [\(5\)](#). 3 % of 1 900 may constitute 57 lives saved per year.

Each year, around 12 000 patients suffer a cerebrovascular stroke [\(6\)](#), and approximately 2 000 of them die [\(7\)](#). The proportion of patients who receive thrombolysis amounts to 21 %, but varies from 5 % to 37 % in different health trusts [\(8\)](#). We may assume that the time from symptom onset to treatment can be reduced if the population is taught how to recognise symptoms of cerebrovascular stroke and how to react.

---

## Life-long learning of first aid

First-aid skills are best maintained by repeated training [\(9\)](#). We therefore believe that a national strategy for life-long training in first aid ought to be developed, where the training starts in the pre-school institutions and is followed up through school, higher education, leisure activities, working life and into the retirement years [\(10\)](#). In addition, special training in life-saving first aid should be provided to health personnel, to broad target groups through the sports and voluntary organisations, and targeted to groups that perhaps have no previous training, for example newly arrived immigrants.

Providing first aid to someone in a life-threatening condition can be a tough experience [\(11\)](#), and many patients die even though they have received high-quality first aid. We are therefore seeking to establish a system through which first-aiders can contact the health services, both in order to provide feedback about their experiences and to take care of those individuals who need further follow-up.

---

## First responders

First responders are personnel who have undergone standardised training in life-saving first aid and have access to simple medical equipment, and are thus able to provide life-saving first aid before the first professional healthcare personnel arrive. First responders should be alerted in situations where their contribution is likely to provide a health gain [\(12\)](#). First responders may include

personnel from the fire and rescue services, the police, the primary health services (e.g. home nursing services) or voluntary organisations such as the Red Cross, Norwegian People's Aid or the Norwegian Society for Sea Rescue.

---

## Better EMCC

Interaction between the caller and the operator at the EMCC (Emergency Medical Communication Centre, 113) is crucial in emergency situations. Most people know that they can call the EMCC, but not everybody is aware that they can receive guidance and assistance regarding what can be done while help is on the way, for example initiating CPR [\(1\)](#). The role of the EMCC is to provide reassurance to the population in the knowledge that it is always available to help. Its motto, 'Call 113 – and you are no longer alone', expresses the same idea. Knowledge of the medical emergency number 113 must be incorporated into all training in first aid, including the way in which the EMCC takes charge and instructs all callers. Participants in first aid training courses should practise how to save lives in collaboration with the EMCC.

*«The population is an especially crucial actor in the medical emergency chain»*

Since the EMCC operator is the professional partner in the first-aid team, the operators need to receive more training in the role of team leader. A national standard for basic and maintenance training of EMCC operators ought to be developed [\(13\)](#).

Publicly available defibrillators have become visible to the EMCC with the aid of registration at the [www.113.no](http://www.113.no) website, and the authorities are currently seeking to establish automatic positioning of callers on a map in the EMCC.

To provide better help to time-critical incidents, it is crucial to have quality registries of relevant conditions, such as the Norwegian Cardiac Arrest Registry, the Norwegian Myocardial Infarction Registry, the Norwegian Stroke Registry and the Norwegian National Trauma Registry. These registries should collect data on the early identification and treatment of patients. Measures initiated by both the caller and the EMCC should be systematically evaluated with a view to improving the training of the general public as well as that of the EMCC operators.

---

## Saving lives together

The 'Saving lives together' programme focuses on the initial links in the medical emergency chain [\(14, 15\)](#), especially on training and implementation. We emphasise collaboration across agencies and organisations to achieve coordination between the different partners. The authorities help ensure that the content of the training prioritises interventions that will save more lives.

The challenge will be to maintain the focus and the quality once the programme is completed. The measures will thus need to be incorporated into the practices of existing organisations.

Until now, efforts have included the introduction of first-aid training in pre-schools, primary, lower and upper secondary schools and among older people. A standard for first aiders is being prepared and a website with information to the public about first aid is being developed. EMCCs all over the country have initiated simulation training, systematic review of audio logs and registration of call data for medical quality registries. Many of these projects have received financial support from the Gjensidige Foundation, which has granted more than NOK 60 million to various projects under the programme to date. Grants remain available for training purposes and input from further target groups, and we would like to systematise training for workplaces, sports and recently arrived immigrants.

The key objective now is to maintain and improve the structures that help ensure that patients with time-critical conditions meet competent and confident first aiders, with good support from the EMCC and technology, irrespective of where the patient is located or who is the closest to provide help.

---

## LITERATURE

1. Mathiesen WT, Birkenes TS, Lund H et al. Public knowledge and expectations about dispatcher assistance in out-of-hospital cardiac arrest. *J Adv Nurs* 2019; 75: 783–92. [PubMed][CrossRef]
2. Wissenberg M, Lippert FK, Folke F et al. Association of national initiatives to improve cardiac arrest management with rates of bystander intervention and patient survival after out-of-hospital cardiac arrest. *JAMA* 2013; 310: 1377–84. [PubMed][CrossRef]
3. Tjelmeland IBM, Kramer-Johansen J, Nilsen JE et al. Årsrapport for 2017 med plan for forbedringstiltak. Norsk hjertestansregister, 2018. [https://www.kvalitetsregistre.no/sites/default/files/8\\_arsrapport\\_2017\\_norsk\\_hjertestansregister\\_0.pdf](https://www.kvalitetsregistre.no/sites/default/files/8_arsrapport_2017_norsk_hjertestansregister_0.pdf) Lest 3.12.2018.
4. Hansen SM, Ringgren KB, Wissenberg M et al. Hjertestop uden for hospital i Danmark. Dansk Hjertestopregister, 2018. [http://genoplivning.dk/wp-content/uploads/2018/04/Dansk\\_Hjertestopregister2016.pdf](http://genoplivning.dk/wp-content/uploads/2018/04/Dansk_Hjertestopregister2016.pdf) Lest 19.12.2018.
5. Tannvik TD, Bakke HK, Wisborg T. A systematic literature review on first aid provided by laypeople to trauma victims. *Acta Anaesthesiol Scand* 2012; 56: 1222–7. [PubMed][CrossRef]
6. Hjerter- og karregisteret. Rapport for 2015. Oslo: Folkehelseinstituttet, 2016. [https://www.fhi.no/globalassets/dokumenterfiler/hjertekarrapport-arsrapport-for-2015\\_010716.pdf](https://www.fhi.no/globalassets/dokumenterfiler/hjertekarrapport-arsrapport-for-2015_010716.pdf) Lest 5.9.2017.

7. Folkehelseinstituttet. Norgeshelsa statistikkbank: Dødsårsaker, nøkkeltall. <http://www.norgeshelsa.no/norgeshelsa/> Lest 26.9.2017.
  8. Fjærtøft H, Indredavik B, Mørch B et al. Årsrapport 2017 – Med plan for forbedringstiltak. Trondheim: Norsk hjerneslagregister, 2018. [https://www.kvalitetsregistre.no/sites/default/files/1\\_arsrapport\\_2017\\_norsk\\_hjerneslagregister.pdf](https://www.kvalitetsregistre.no/sites/default/files/1_arsrapport_2017_norsk_hjerneslagregister.pdf) Lest 4.12.2018.
  9. Celenza T, Gennat HC, O'Brien D et al. Community competence in cardiopulmonary resuscitation. *Resuscitation* 2002; 55: 157–65. [PubMed] [CrossRef]
  10. Norges offentlige utredninger. Først og fremst. Et helhetlig system for håndtering av akutte sykdommer og skader utenfor sykehus. 2015:17. <https://www.regjeringen.no/contentassets/477c27aa89d645e09ece350eaf93fedf/no/pdfs/nou20152015001700odddpdfs.pdf> Lest 19.12.2018.
  11. Mathiesen WT, Bjørshol CA, Braut GS et al. Reactions and coping strategies in lay rescuers who have provided CPR to out-of-hospital cardiac arrest victims: a qualitative study. *BMJ Open* 2016; 6: e010671. [PubMed] [CrossRef]
  12. Rundskriv I-5/15. Akuttmedisinforskriften. Med merknader til enkelte bestemmelser. Oslo: Helse- og omsorgsdepartementet, 2015.
  13. Anbefalinger fra delprosjekt 113 – Sammen redder vi liv. Oslo: Helsedirektoratet, 2018. <https://www.helsedirektoratet.no/rapporter/sammen-redder-vi-liv-strategidokument/Saving%20lives%20together%20-%20Recommendations%20from%20project%20113.pdf?download=true> Lest 15.9.2018.
  14. Nasjonal dugnad for å redde liv ved hjertestans og andre akuttmedisinske tilstander utenfor sykehus. Strategi IS-2702. Oslo: Helsedirektoratet, 2017. <https://www.helsedirektoratet.no/rapporter/sammen-redder-vi-liv-strategidokument/Saving%20lives%20together%20-%20Strategy.pdf?download=true> Lest 16.12.2017.
  15. Faktagrunnlag – Sammen redder vi liv. Oslo: Helsedirektoratet, 2018. [http://www.helsedirektoratet.no/rapporter/sammen-redder-vi-liv-strategidokument/Sammen%2oredder%20vi%20liv%20-%20faktagrunnlag.pdf/\\_/attachment/download/40cob34f-3c7e-4ca3-9dc1-65bd4e44ad89:56479c73b871f2f67411b6ba24d8acbce9c5df69/Sammen%2oredder%20vi%20liv%20-%20faktagrunnlag.pdf](http://www.helsedirektoratet.no/rapporter/sammen-redder-vi-liv-strategidokument/Sammen%2oredder%20vi%20liv%20-%20faktagrunnlag.pdf/_/attachment/download/40cob34f-3c7e-4ca3-9dc1-65bd4e44ad89:56479c73b871f2f67411b6ba24d8acbce9c5df69/Sammen%2oredder%20vi%20liv%20-%20faktagrunnlag.pdf) Lest 19.12.2018.
- 

Publisert: 9 September 2019. Tidsskr Nor Legeforen. DOI: 10.4045/tidsskr.19.0128  
Received 9.2.2019, first revision submitted 3.5.2019, accepted 10.5.2019.  
Copyright: © Tidsskriftet 2026 Downloaded from tidsskriftet.no 14 February 2026.