
Patient visits to an emergency department in anticipation of the COVID-19 pandemic

SHORT REPORT

LARS PETTER BJØRNSSEN

E-mail: lars.p.b.w.bjornsen@ntnu.no

Emergency Department

Division of Emergency Medicine and Prehospital Care

St. Olav's Hospital, Trondheim University Hospital
and

Department of Circulation and Medical Imaging

NTNU – Norwegian University of Science and Technology

Lars Petter Bjørnsen is a specialist in emergency medicine, a senior
consultant and an associate professor.

The author has completed the ICMJE form and declares no conflicts of
interest.

LARS EIDE NÆSS-PLEYM

Emergency Department and

Department of Emergency Medicine

Division of Emergency Medicine and Prehospital Care

St. Olav's Hospital, Trondheim University Hospital

Lars Eide Næss-Pleym is an ICT adviser and doctoral research fellow.

The author has completed the ICMJE form and declares no conflicts of
interest.

JOSTEIN DALE

Department of Emergency Medicine

Division of Emergency Medicine and Prehospital Care

St. Olav's Hospital, Trondheim University Hospital

Jostein Dale is a senior consultant and head of department.

The author has completed the ICMJE form and declares no conflicts of
interest.

Emergency Department

Division of Emergency Medicine and Prehospital Care

St. Olav's Hospital, Trondheim University Hospital

and

Department of Circulation and Medical Imaging

NTNU – Norwegian University of Science and Technology

Lars Erik Laugsand is a senior consultant and assistant head of division.

The author has completed the ICMJE form and declares no conflicts of interest.

BACKGROUND

The Emergency Department in Trondheim has prepared for a large influx of patients infected with the SARS-CoV-2 virus. We conducted a study comparing patients in the Emergency Department in the first weeks of the pandemic in Norway (weeks 11 and 12) with the average number of patient visits.

MATERIAL AND METHOD

Data from patients at the Emergency Department of St Olav's Hospital in the period 6 January 2020–22 March 2020 were retrieved from the Emergency Department's database. Logistical patient data concerning patient numbers, chief complaints, length of stay in the Emergency Department, acuity level, isolation status, and treatment level were analysed.

RESULTS

In week 12, 331 patients were referred to the Emergency Department, a reduction of 39 % compared with the average of 541 patients in weeks 2–10. There was a general reduction in all patient groups, but particularly those discharged from the Emergency Department. In week 12 there were 56 more patients isolated with suspected/potentially infectious disease (187 %) compared with the average for weeks 2–10, and these patients spent almost two hours longer in the Emergency Department than other patients.

INTERPRETATION

There was a reduction in patient visits to the Emergency Department in the first weeks of the pandemic. The percentage of patients isolated for infection control increased, and the time spent in the Emergency Department for these patients was greater than for other patients. The reduction in the inflow of patients is expected to be temporary, and the Emergency Department at St Olav's Hospital expects a large influx of patients with suspected COVID-19 disease.

Main findings

In the course of the first weeks of the COVID-19 pandemic in Norway there was a general reduction in the number of patients referred to the Emergency Department.

The reduction was greatest among patients with presumed low acuity levels and patients normally discharged directly from the Emergency Department.

The number of patients isolated because of suspected infection increased, and these patients spent substantially longer in the Emergency Department.

The current pandemic caused by the SARS-CoV-2 virus was predicted to reach Norway shortly after the outbreak in China [\(1\)](#). Following the government decision to virtually shut Norway down on 12 March (week 11), acute admissions departments have prepared for a large influx of infected patients. The Emergency Department at St Olav's Hospital, Trondheim University Hospital is preparing for worst-case scenarios, but is now experiencing the calm before the storm. No Emergency Department health personnel are in quarantine, but as of 14 April 2020 a total of 15 members of the staff of St Olav's Hospital have COVID-19 disease and 83 are in quarantine or at home with respiratory tract symptoms [\(2\)](#)

In recent years, the Emergency Department at St Olav's Hospital has seen a gradual increase in patient visits [\(3\)](#). The patient population consists primarily of patients aged over 15 years. Acuity level is in all cases determined using the Rapid Emergency Triage and Treatment System (RETTS) [\(4\)](#) according to which patients are categorised as red, orange, yellow or green depending on assumed severity. Red patients must be seen immediately by a doctor, while green patients can wait if more urgent cases come at the same time. Children and adults presenting with problems in the fields of obstetrics and gynaecology, ophthalmology and ear, nose and throat are as a general rule received at separate receptions or clinics [\(3\)](#).

In this study we wanted to compare the inflow of patients in weeks 11 and 12, when St Olav's Hospital stepped up its preparations for the pandemic, with the average for the previous weeks (weeks 2–10).

Material and method

Data from all patients who attended the Emergency Department at St Olav's Hospital in the period 6 January 2020 to 22 March 2020 (weeks 2–10) were retrieved from the Emergency Department's database (Akuttdatabasen, Helse-Vest ICT, version 1.5.5., Stavanger). We chose this period because we assume that it is representative of a stable flow of patients unaffected by public holidays. Logistical data such as patient numbers, medical issue, time spent in the Emergency Department, urgency, infection status and treatment level, were

imported and processed using the Statistical Package for the Social Sciences (SPSS) version 25. Relevant data were then transferred to Microsoft Excel 2016 for graphical presentations. The study was approved by the data protection official (ESA no. 16/9114) and submitted to the Regional Committee for Medical and Health Research Ethics (REK 2016/1813).

Results

In Weeks 11 and 12, 426 and 331 patients, respectively, were referred to the Emergency Department, compared with the average of 541 patients in weeks 2–10. This is equivalent to a reduction of 115 patients (21 %) in week 11 and 210 patients (39 %) in week 12 (Fig. 1). The reduction was clear already in the middle of week 11 (Figure 2).




Figure 1 Overview of the total number of patients seen weekly in the Emergency Department at St Olav's Hospital in the first quarter of 2020 and the weekly number of patients isolated in the Emergency Department in the same period because of suspected/potentially infectious disease.




Figure 2 Number of inpatients in the Emergency Department at St Olav's Hospital each hour on the hour in weeks 11 and 12 in 2020 compared with average visits in weeks 2–10 in 2020.

In week 12 there was a general reduction in acute patients in all medical areas. The reductions in patients in the areas medicine, surgery, orthopaedics and neurology were 98 (35 %), 49 (39 %), 16 (41 %) and 16 (40 %), respectively, compared with the averages for weeks 2–10. For urgency, there were reductions in red, orange, yellow and green of 23 (39 %), 84 (41 %), 87 (36 %) and 13 (48 %) patients. The number of outpatients treated at the clinic was reduced by 76 (47 %). There was no change in the number of patients admitted to the Intensive Care Department or Observation Unit (data not shown).

There was a marked increase in weeks 11 and 12 in patients who were isolated in the Emergency Department because of suspected/potentially infectious disease (Fig. 1). The average time spent by patients in the Emergency Department was 3 hours 14 minutes in weeks 2–10, 2 hours 52 minutes in week 11 and 3 hours 29 minutes in week 12. The time spent by patients in infection isolation was 5 hours 1 minute in week 12, despite the fact that the occupancy rate at the hospital in week 12 was 65 % (internal data from St Olav's Hospital, PAS-rapporten Korpas database).

Discussion

We found a significant reduction in the number of patients referred to the Emergency Department early in the epidemic. A similar situation was reported during the first two weeks of the epidemic in Italy [\(5\)](#) and in China, where a sharp reduction was seen in the number of patients seeking emergency dental treatment [\(6\)](#). However, there are no studies that describe this trend in more detail. We saw a general reduction in the number of patients coming to the Emergency Department from week 11 already, without any medical disciplines standing out. Other European countries such as Belgium, France and Germany

refer to unpublished data on a general reduction in the number of patients coming to the Emergency Department, particularly patients with conditions assumed to be less serious (7).

Our study revealed a similar reduction in the number of patients with low urgency and patients whose treatment would normally be fully dealt with in the Emergency Department. In Norway this may be because the Norwegian primary health service has largely dealt with these issues itself, or because the threshold for referring patients to the Emergency Department has risen now in a time of crisis. The primary health service makes sound clinical judgements that limit the inflow of patients. The fact that the inflow of patients is almost unchanged at the weekend, when GP offices are closed, provides support for this statement (Fig. 2).

An explanatory model for the pronounced fall that is greater cause for concern is that patients are delaying contacting their GP or phoning the emergency number for fear of becoming infected or of being an unnecessary burden on the health service. This fear is shared by emergency doctors in Belgium, the UK and France, among others. From Germany there are reports of a reduction in stroke and heart attack cases in the Emergency Department (7), and there have been similar anecdotal reports in Trondheim, Norway. Possibly in order to avoid being a burden on the health system, or for fear of infection, patients have suffered both stroke and heart attacks without seeking help within treatment time window.

It is crucial that the primary health service acts as gatekeeper and filter for the specialist health service. At the same time, the message must be communicated to the public that people *must* seek medical assistance as early as possible in the event of symptoms of what may be a serious illness.

The aim during the pandemic is for all patients to have safe, high quality assessment and treatment at the Emergency Department, no matter what the situation. This is challenging, as assessment and treatment of isolated patients takes longer and requires more resources. A rapid test for COVID-19 in the Emergency Department would probably reduce clarification time in the Emergency Department. The greatest potential challenge, however, is to have a sufficient number of health personnel (2). Although no Emergency Department staff are in quarantine at present, there is a high risk of health personnel becoming unavailable. It is the calm before the storm in Trondheim, and the Emergency Department is preparing for a large influx of patients with suspected COVID-19.

LITERATURE

1. Aavitsland P. The coronavirus epidemic will reach Norway. Tidsskr Nor Legeforen 2020; 140. doi: 10.4045/tidsskr.20.0077. [PubMed][CrossRef]
2. St. Olavs hospital. Beredskap. <https://stolav.no/beredskap> Accessed 13.4.2020.
3. Bjørnsen LP, Uleberg O, Dale J. Patient visits to the emergency department at a Norwegian university hospital: variations in patient gender and age,

timing of visits, and patient acuity. *Emerg Med J* 2013; 30: 462–6. [PubMed]
[CrossRef]

4. Widgren BR, Jourak M. Medical Emergency Triage and Treatment System (METTS): a new protocol in primary triage and secondary priority decision in emergency medicine. *J Emerg Med* 2011; 40: 623–8. [PubMed][CrossRef]

5. Coen D, Paolillo C, Cavazza M et al. Changing emergency department and hospital organization in response to a changing epidemic. *Emergency Care Journal* 2020; 16: 8969.

6. Guo H, Zhou Y, Liu X et al. The impact of the COVID-19 epidemic on the utilization of emergency dental services. *J Dent Sci* 2020; 15. doi: 10.1016/j.jds.2020.02.002. [PubMed][CrossRef]

7. The European Society for Emergency Medicine. COVID-19 Webinar: a unique insight in highly impacted countries. Sharing experiences of high COVID-19 impact countries. https://academy.eusem.org/eusem/2020/covid-19/290502/session.speakers.covid-19.webinar.a.unique.insight.in.highly.impactted.countries.html?f=menu=8*browseby=8*sortby=2*label=19798 Accessed 13.4.2020.

Publisert: 24 April 2020. *Tidsskr Nor Legeforen*. DOI: 10.4045/tidsskr.20.0277

Received 29.3.2020, first revision submitted 14.4.2020, accepted 21.4.2020.

Published under open access CC BY-ND. Downloaded from tidsskriftet.no 28 December 2025.