
Gas in soft tissue

IMAGES IN MEDICINE

JAKOB LILLEMOEN DRIVENES

jakobdrivenes@hotmail.com

Division of Surgery

Vestfold Hospital Trust

Jakob Lillemoen Drivenes, specialty registrar (part 1).

The author has completed the ICMJE form and declares no conflicts of interest.

BJØRN ÅSHEIM HANSEN

Department of Infectious Diseases

Vestfold Hospital Trust

Bjørn Åsheim Hansen, senior consultant.

The author has completed the ICMJE form and declares no conflicts of interest.

MILOS STOJANOVIC

Department of Radiology

Vestfold Hospital Trust

Milos Stojanovic MD PhD, senior consultant.

The author has completed the ICMJE form and declares no conflicts of interest.

HENRIK JOHAN STØREN

Department of Orthopaedics

Vestfold Hospital Trust

Henrik Johan Støren, senior consultant.

The author has completed the ICMJE form and declares no conflicts of interest.

ÅSHILD MARVIK

Department of Microbiology

Vestfold Hospital Trust

Åshild Marvik, senior consultant.

The author has completed the ICMJE form and declares no conflicts of interest.



This X-ray image of the left ankle shows gas in the soft tissue around the heel. The patient, a man in his seventies, was admitted to the emergency department with fever and pain in the left foot. He was known to have colon cancer with spread to the peritoneum and lungs. Four years prior to this admission, he had undergone right hemicolectomy, but had declined any further curative treatment. Three days prior to admission, he developed pain in the left ankle and, according to his spouse, he became confused and his condition declined further. There was no information about trauma, recent wounds or rash.

On arrival, the patient was confused (Glasgow Coma Scale score of 14), febrile (38.8 °C rectal temperature) and tachycardic (140 bpm), but he had normal blood pressure (129/68 mmHg). Blood test results showed CRP 166 mg/L (reference range <5) and leukocytes $20.3 \times 10^9/L$ ($3.5-10 \times 10^9/L$). The findings of examination of the left ankle were unremarkable, but an intact blister over the heel was noticed, as well as an erythematous rash over the medial side of the left calf. In addition to intravenous fluid administration, treatment for suspected cellulitis was initiated with cloxacillin 2 g four times daily and intravenous gentamicin 5 mg/kg body weight once daily.

The following day, there was growth of *Clostridium septicum* in two anaerobic blood culture bottles. At this time, the blister over his heel spontaneously ruptured, releasing extremely foul-smelling fluid. There was no definite palpable crepitus. X-ray of the left ankle revealed soft tissue emphysema around the heel. Antibiotic treatment was switched to intravenous piperacillin/tazobactam 4 g/0.5 g four times daily. One and a half days after admission, the patient underwent supramalleolar guillotine amputation as a life-saving procedure under general anaesthesia, and vacuum-assisted closure therapy was initiated. One week later, definitive below-knee amputation of the left limb was performed with primary closure. After a further week of observation, the condition of the amputation stump was unremarkable. The patient was discharged for a short-term admission in a municipal institution and referred for rehabilitation to have a prosthesis fitted.

Clostridia are gram-positive, anaerobic, spore-forming bacteria, which are widespread in the environment, as well as in the gastrointestinal tract of healthy humans and animals. Clostridial myonecrosis, or gas gangrene, has a high mortality rate, and early surgical intervention and antibiotic treatment are essential (1, 2). There are two types of gas gangrene – traumatic and spontaneous. *C. perfringens* is the most common species in traumatic gas gangrene, while *C. septicum* is the predominant agent in spontaneous gas gangrene (2). Underlying disease or an iatrogenic procedure in the gastrointestinal tract is a prerequisite for the haematogenous spread of bacteria to the muscle system. Infections with *C. septicum* are associated with malignancy, particularly in the gastrointestinal tract, and this diagnosis should therefore be kept in mind when patients with colon cancer develop sepsis and soft tissue disorders (1, 2).

In this case, the blood culture findings were crucial, but palpable crepitus is considered to be the most important clinical finding for early diagnosis.

The patient has consented to the publication of the article.

The article has been peer-reviewed.

REFERENCES

1. Bernardshaw SV, Røkke O. Clostridium septicum-infeksjon og kreft. Tidsskr Nor Lægeforen 2001; 121: 2381. [PubMed]
2. Srivastava I, Aldape MJ, Bryant AE et al. Spontaneous C. septicum gas gangrene: A literature review. Anaerobe 2017; 48: 165–71. [PubMed]

[CrossRef]

Publisert: 9 February 2023. Tidsskr Nor Legeforen. DOI: 10.4045/tidsskr.22.0686

Received 26.10.2022, first revision submitted 8.12.2022, accepted 2.1.2023.

Copyright: © Tidsskriftet 2026 Downloaded from tidsskriftet.no 21 March 2026.