

Infection prevention practices can cause hand eczema

FROM THE SPECIALTIES

INGVILL SANDVEN

ingvsa@ous-hf.no

Ingvill Sandven, specialist in occupational medicine and senior consultant at Occupational Medicine and Environmental Health, Oslo University Hospital.

The author has completed the ICMJE form and declares no conflicts of interest.

HILDE HEIRO

Hilde Heiro, specialist in occupational medicine and senior consultant at Occupational Medicine and Environmental Health, Oslo University Hospital.

The author has completed the ICMJE form and declares no conflicts of interest.

EGIL LINGAAS

Egil Lingaas, specialist in medical microbiology and head senior consultant at the Department of Infection Prevention, Oslo University Hospital.

The author has completed the ICMJE form and declares no conflicts of interest.

TERESA LØVOLD BERENTS

Teresa Løvold Berents, PhD, specialist in dermatovenereology, qualified in allergology, and senior consultant at the Department of Dermatology and Regional Centre for Asthma, Allergy and Hypersensitivity (RAAO), Oslo University Hospital.

The author has completed the ICMJE form and declares no conflicts of interest.

Hand eczema prevention should be covered in guidelines on hand hygiene. A new training film can be helpful.

Good hand hygiene is essential for preventing the spread of infection in healthcare settings. However, it can lead to hand eczema, which is not only a problem for individual employees but also increases the risk of infection transmission.

Hand eczema is twice as common among healthcare personnel as in the general population. Studies from several countries found a further increase in the incidence of hand eczema among healthcare personnel during the COVID-19 pandemic, which is attributed to the more stringent hand hygiene requirements in place [\(1, 2\)](#).

To prevent work-related hand eczema, it is recommended that employees in high-risk occupations, such as healthcare personnel, are trained in how to care for their skin and reduce risk factors at an early stage, for example through the correct use of gloves and hand sanitiser [\(3\)](#).

There is a clear correlation between poor hand hygiene and an increased risk of infection. All healthcare institutions are therefore required to have written guidelines for hand hygiene. Dermatological problems resulting from hygiene practices are known to be one of the factors leading to non-compliance with these measures.

The guidelines often recommend practising hand hygiene before putting on non-sterile disposable gloves. This is justified by the possibility of contaminating the outside of the gloves when putting them on, and contaminating other gloves in the same box. However, this is not well documented. On the contrary, several studies indicate that this is not a significant problem [\(4–6\)](#). If the skin is moist before donning gloves, this can increase the risk of hand eczema. We therefore think that this recommendation should be removed from the guidelines.

Efforts to prevent hand eczema should be viewed in conjunction with hand hygiene guidelines and training, as maintaining healthy skin is vital for adhering to these guidelines and also for effective infection control.



Screen shot from training film

At Oslo University Hospital, the Department of Infection Prevention, the Department of Dermatology, and Environmental and Occupational Medicine have jointly produced a short training film giving advice on how to avoid hand eczema (7). The film contains information that we believe is useful for all staff working in a clinical setting, and serves as a practical tool that managers can use in staff training.

REFERENCES

1. Vindenes HK, Bertelsen RJ, Lygre SHL et al. Changes in Infection Prevention Practices and Occurrence of Skin Symptoms among Healthcare Workers, Cleaners and Day-care Workers in Norway during the COVID-19 Pandemic. *Acta Derm Venereol* 2023; 103. doi: 10.2340/actadv.v103.3420. [PubMed][CrossRef]
2. Erdem Y, Altunay IK, Aksu Çerman A et al. The risk of hand eczema in healthcare workers during the COVID-19 pandemic: Do we need specific attention or prevention strategies? *Contact Dermat* 2020; 83: 422–3. [PubMed][CrossRef]
3. Thyssen JP, Schuttelaar MLA, Alfonso JH et al. Guidelines for diagnosis, prevention, and treatment of hand eczema. *Contact Dermat* 2022; 86: 357–78. [PubMed][CrossRef]
4. Lingaas E, Fagernes M. Development of a method to measure bacterial transfer from hands. *J Hosp Infect* 2009; 72: 43–9. [PubMed][CrossRef]
5. Thom KA, Rock C, Robinson GL et al. Direct gloving vs hand hygiene before donning gloves in adherence to hospital Infection control Practices. *JAMA Netw Open* 2023; 6. doi: 10.1001/jamanetworkopen.2023.36758. [PubMed][CrossRef]
6. Rock C, Harris AD, Reich NG et al. Is hand hygiene before putting on nonsterile gloves in the intensive care unit a waste of health care worker time?—a randomized controlled trial. *Am J Infect Control* 2013; 41: 994–6. [PubMed][CrossRef]
7. Vestre Viken HF. Arbeidsrelatert håndeksem. Vimeo 14.4.2023. <https://vimeo.com/817593664> Accessed 4.12.2023.

Publisert: 8 January 2024. Tidsskr Nor Legeforen. DOI: 10.4045/tidsskr.23.0759

Received 6.11.2023, first revision submitted 24.11.2023, accepted 4.12.2023.

Copyright: © Tidsskriftet 2026 Downloaded from tidsskriftet.no 26 March 2026.