

Foreign body aspiration in a child

SHORT CASE REPORT

LINE NORMAN KVENSHAGEN

line.norman.kvenshagen@so-hf.no

Department of Paediatric and Adolescent Medicine
Østfold Hospital Kalnes

Line Norman Kvenshagen, specialist in paediatrics and senior consultant
The author has completed the ICMJE form and declares no conflicts of interest.

GURO GRINDHEIM

Department of Anaesthesia and Intensive Care Medicine

Oslo University Hospital, Rikshospitalet

Guro Grindheim PhD, specialist in anaesthesiology, senior consultant
and head of unit

The author has completed the ICMJE form and declares no conflicts of interest.

JENS KRISTIAN STENE-JOHANSEN

Department of Anaesthesia and Intensive Care Medicine

Østfold Hospital Kalnes

Jens Kristian Stene-Johansen, specialist in anaesthesiology and senior
consultant

The author has completed the ICMJE form and declares no conflicts of interest.

EIRIK FRANER

Department of Anaesthesia and Intensive Care Medicine

Østfold Hospital Kalnes

Eirik Franer, specialist in anaesthesiology and senior consultant

The author has completed the ICMJE form and declares no conflicts of interest.

CHRISTIAN RIDDERVOLD KAHRS

Department of Paediatric and Adolescent Medicine

Østfold Hospital Kalnes

Christian Riddervold Kahrs PhD, specialist in paediatric medicine and senior consultant

The author has completed the ICMJE form and declares no conflicts of interest.

KRISTINA SVENNERHOLM

Department of Anaesthesia and Intensive Care Medicine

Østfold Hospital Kalnes

and

Department of Anaesthesia and Intensive Care Medicine

Sahlgrenska University Hospital

Gothenburg

Kristina Svennerholm, senior consultant

The author has completed the ICMJE form and declares no conflicts of interest.

MARTE KRISTINE TVEIT MYHRE

Department of Radiology

Østfold Hospital Kalnes

Marte Kristine Tveit Myhre, senior consultant

The author has completed the ICMJE form and declares no conflicts of interest.

Background

Aspiration of a foreign body can lead to a fatal outcome, especially in children younger than 3 years of age.

Case presentation

A healthy 2-year-old child was admitted to the emergency unit due to cough and respiratory distress. They were admitted to the paediatric ward with a tentative diagnosis of pneumonia. After some hours, the child deteriorated, showing severe respiratory compromise. Non-invasive respiratory support and inhalation therapy were ineffective. The child was intubated, but mechanical ventilation was not feasible due to hypercapnia. Bronchoscopy was carried out and several foreign bodies were removed. Visual examination revealed fragments of almonds. The child was discharged from hospital after 12 days.

Interpretation

The child underwent aspiration. After intubation, the almonds may have caused a ball-valve effect, leading to the life-threatening situation. It is important to consider a foreign body as a potential cause of abnormal respiratory symptoms in children.

This case report describes airway aspiration of a foreign body in a child. The child initially presented with mild respiratory symptoms but rapidly progressed to respiratory failure and life-threatening complications.

A two-year-old child was referred to the emergency department by their general practitioner (GP) due to respiratory symptoms. The father reported that the child had been coughing for two weeks without any additional symptoms. However, two days before admission, the child's condition deteriorated, they were less active and the cough worsened. The GP suspected a viral respiratory infection and referred the child to the paediatric department at the hospital.

The duty paediatricians examined the child upon arrival. The child was alert but pale and fatigued. Breathing was laboured with intercostal and subcostal retractions. On auscultation, wheezing and crackles were heard from the lungs. The respiratory rate was 80 - 90 breaths per minute (reference range 25 - 35). Heart auscultation findings were normal, and the pulse was 180 beats per minute (90 - 130). Oxygen saturation was 74 % on room air and 95 % with 7 litres of oxygen on a mask. The child was afebrile.

The blood tests showed CRP 214 mg/L (<5), white blood cells $16.9 \times 10^9/L$ ($3.5-14.0 \times 10^9/L$), capillary pH 7.42 (7.35-7.45), and pCO_2 4.2 kPa (4.7-5.9).

The condition was interpreted as pneumonia, and the patient was admitted to the paediatric ward. A total of 315 mg of penicillin was administered intravenously, and high-flow nasal cannula therapy was started at a rate of 15-25 litres per minute and 40 % oxygen supplementation. Inhalations with 3 ml NaCl and 2.5 mg of salbutamol had an uncertain effect. Chest X-ray showed perihilar opacities (Figure 1). After five hours, the child appeared more fatigued but was still aware of their surroundings. Breathing entailed wheezing with prolonged expiration. The duty paediatrician noted diminished breath sounds on the right side.




Figure 1 Chest X-ray on admission to hospital showed perihilar opacities.

A total of 48 mg of intravenous hydrocortisone was administered, along with more frequent inhalations of NaCl and salbutamol in the same dose as before. A capillary blood gas test at this time showed a pH of 7.29 and PCO_2 of 6.7 kPa. The child was transferred to the intensive care unit. Swelling in the neck was observed. A new chest X-ray showed pneumomediastinum and soft tissue emphysema in the neck and left side of the chest.

The child's condition had now deteriorated to the point of extreme fatigue and severe respiratory compromise, leading to the joint decision by the paediatricians and anaesthesiologists that intubation was necessary. The child was premedicated with ketamine, fentanyl and rocuronium bromide and was intubated by an anaesthesiologist 11 hours after admission. Immediately after intubation, condensation was observed in the tube, and the response from end-tidal CO_2 measurement indicated correct tube placement. However, there were minimal chest movements and weak, bilateral breath sounds. The child was placed on a ventilator with ketamine and propofol infusion, along with bolus doses of fentanyl and cisatracurium, but required unacceptably high pressures and volumes. As a result, the patient was manually ventilated with 100 % oxygen supplementation. Nevertheless, satisfactory ventilation or oxygenation could not be achieved. Oxygen saturation (SpO_2) remained around 70 %. Arterial blood gas at this point showed a pH of 6.8, PO_2 16.4 kPa and PCO_2 24 kPa.

The university hospital was contacted regarding treatment with extracorporeal membrane oxygenation (ECMO), and it was decided to send the ECMO team to the local hospital.

While awaiting the team's arrival, inhaled 20 ppm nitric oxide was administered (1). A new inhalation of 2.5 mg of salbutamol, along with 120 mg of intravenous theophylline and 625 mg of magnesium sulphate was also administered, but no improvement was observed in the patient. Circulatory stability was maintained with an infusion of 0.1 µg/kg/min adrenaline, but there was a life-threatening ventilation issue. It was decided to insert bilateral chest drains due to vital indications, as minimal breath sounds were heard and there were clinical and radiological signs of worsening emphysema. A subsequent chest X-ray showed increasing opacity, pneumomediastinum and emphysema (Figure 2). The antibiotic treatment was changed to a combination of 625 mg of cefotaxime and 120 mg of clindamycin intravenously.




Figure 2 X-ray taken after intubation and chest drains on both sides. There was no pneumothorax, but increased opacity. Pronounced pneumomediastinum and soft tissue emphysema were observed over the face, neck and left hemithorax.

The ECMO team arrived at the local hospital four hours after intubation. Bronchoscopy revealed a significant amount of mucus and swollen mucous membranes. Adequate ventilation was subsequently achieved, allowing transport to the university hospital without ECMO treatment. A pulmonologist at the university hospital performed a bronchoscopy under anaesthesia, which

lasted four hours and resulted in the removal of two lodged foreign bodies in the left upper lobe and middle lobe bronchi. These appeared to be fragments of almonds. During a subsequent conversation with the patient's parents, it emerged that the child had eaten granola with almonds at kindergarten two days before admission, the same day the situation became more severe.

The child was extubated after two days in the paediatric intensive care unit. In the following week, the child required high-flow nasal cannula therapy at night, intermittent CPAP treatment during the day and chest physiotherapy. A CT of the chest showed total atelectasis of the right middle lobe and left lower lobe. The child was discharged from the hospital with a plan for inhalation therapy and a subsequent check-up with bronchoscopy.

Discussion

Nuts are the most common cause of foreign body aspiration in children, with the highest incidence between the ages of 1 and 3 years (2, 3). Coughing, wheezing and choking/gagging can be symptoms of foreign body aspiration but may also indicate respiratory infections (3, 4). Fever is typically associated with infections, which our patient did not have. However, CRP was elevated, which is not necessarily the case in foreign body aspiration.

Most children who are evaluated for suspected foreign body aspiration undergo a chest X-ray (> 80 %), and 30–50 % of these are reported to have normal findings (3, 5). Our patient's X-ray showed signs of infection but no suspicion of a foreign body. Given the two-week history of coughing, it is possible, in hindsight, that the child may also have had a mild viral respiratory infection.

The patient may have had partially obstructed airways upon arrival. It is possible that the positive pressure ventilation after intubation displaced the foreign bodies, leading to almost complete airway obstruction and a ball-valve effect, causing the persistently high pCO₂ level (6).

Foreign body aspiration can have a fatal outcome (7, 8), and the condition of the child in our case report was life threatening. Bronchoscopy is necessary to remove the foreign body in cases of aspiration (4, 9). Foreign body aspiration should always be considered in children with unusual presentations of respiratory symptoms.

The patient's parents have consented to publication of the article.

The article has been peer-reviewed.

REFERENCES

1. Kinsella JP, Abman SH. Inhaled nitric oxide therapy in children. *Paediatr Respir Rev* 2005; 6: 190–8. [PubMed]
2. Molla YD, Mekonnen DC, Beza AD et al. Foreign body aspiration in children at University of Gondar Comprehensive Specialized Hospital, a two

year retrospective study. *Heliyon* 2023; 9. doi: 10.1016/j.heliyon.2023.e21128. [PubMed][CrossRef]

3. Sink JR, Kitsko DJ, Georg MW et al. Predictors of Foreign Body Aspiration in Children. *Otolaryngol Head Neck Surg* 2016; 155: 501–7. [PubMed]
4. Stevens R, Kelsall-Knight L. Clinical assessment and management of children with bronchiolitis. *Nurs Child Young People* 2022; 34: 13–21. [PubMed]
5. Hutchinson KA, Turkdogan S, Nguyen LHP. Foreign body aspiration in children. *CMAJ: Canadian Medical Association journal = journal de l'Association medicale canadienne* 2023; 195: E333.
6. Kent J, Ng C. Foreign body airway obstruction causing a ball valve effect. *JRSM Short Rep* 2013; 4.. [PubMed]
7. Mîndru DE, Păduraru G, Rusu CD et al. Foreign Body Aspiration in Children-Retrospective Study and Management Novelties. *Medicina (Kaunas)* 2023; 59: 1113. [PubMed]
8. Rodríguez H, Passali GC, Gregori D et al. Management of foreign bodies in the airway and oesophagus. *Int J Pediatr Otorhinolaryngol* 2012; 76 (Suppl 1): S84–91. [PubMed][CrossRef]
9. Fidkowski CW, Zheng H, Firth PG. The anesthetic considerations of tracheobronchial foreign bodies in children: a literature review of 12,979 cases. *Anesth Analg* 2010; 111: 1016–25. [PubMed][CrossRef]

Publisert: 26 March 2025. Tidsskr Nor Legeforen. DOI: 10.4045/tidsskr.24.0390

Received 9.7.2024, first revision submitted 14.11.2024, accepted 10.1.2025.

Published under open access CC BY-ND. Downloaded from tidsskriftet.no 14 February 2026.